# Summary of electron response at phi crack

University of Rochester Geumbong Yu, Yeonsei Chung


### Event Selection

- $\bigcirc$  MC: 1 million zewkae (Z $\rightarrow$ ee)
- Data: bhelOd (single high pt electron)
- Require one electron by em object and the other track to reconstruct Z. (tower 1~8)
- No background assumed.
- Central tight electron cut on electron leg, pt > 20 GeV cut instead of et cut
- Cuts for the other track(2x2 Energy used):
  - Opposite charge to electron's, Had/Em < 0.5 E/p < 1.2, pt > 20 GeV, No other track within 0.4 cone

## Track Had/Em & E/p distribution


Blue: track | relative phi | < 0.9

Green: track |relative phi|>=0.9
Data in marker while MC in line



Single tower E


Track momentum



### Z mass after selection cut

#### Monte Carlo


#### Data




### Z mass vs. relative phi in tower

#### Monte Carlo

Data
Energy 4-vec




Momentum 4-vec



### Z mass distribution

Blue: track |relative phi|<0.9 Green: track |relative phi|>=0.9 Data in marker while MC in line

Momentum 4-vec



## E/p vs relative position

- @ Good region required:
  - for E/p vs. rel. phi, |reta|<0.5 used.
  - for E/p vs. rel. eta, |rphi|<0.5 used.
- Relative position?
  - Scaled track hit position on CES face in eta or phi direction of a tower.


$$rphi(reta) = \frac{glob \_phi(eta) - \frac{max\_phi(eta) + min\_phi(eta)}{2}}{\frac{max\_phi(eta) - min\_phi(eta)}{2}}$$

## Track E/p vs. relative phi

#### Monte Carlo

Difference = (data)E/p - (mc)E/p

Data




## Track E/p vs. relative eta

#### Monte Carlo

Difference = (data)E/p - (mc)E/p

Data




Aug 25 2005 Simulation Meeting University of Rochester Geumbong Yu, Yeonsei Chung

## Electron E/p vs. relative phi

#### Monte Carlo

Data




## Electron E/p vs. relative eta


#### Monte Carlo

Difference = (data)E/p - (mc)E/p

Data

(mc)E/p






## Em object E/p vs. relative phi


#### Monte Carlo

Difference = (data)E/p - (mc)E/p

Data

(mc)E/p






## Em object E/p vs. relative eta


#### Monte Carlo

Difference = (data)E/p - (mc)E/p


Data

(mc)E/p






# Good region dependence in E/p vs. emobject phi



# Good region dependence in E/p vs. emobject eta




### Conclusion.


- No background study is done for this study.
- Comparing single tower energy divided by momentum of Monte Carlo with data shows a couple of % difference in central region but more discrepancy in high reta(rphi) region. Emobject shows good agreement between Monte Carlo and data.
- After reconstruction, the Monte Carlo agrees with data well by checking the E/p from emobject. Map or leakage correction for Cal data?

## Backup plots


## Zmass with different E/p cut




# Good region dependence in E/p vs. track rphi




# Good region dependence in E/p vs. track reta



# Good region difference in E/p vs. electron rphi




# Good region dependence in E/p vs. electron reta




### E/p of west and east calorimeter


West & East

Monte Carlo described in line and Data in marker





## Emobject E/p - single E/p



Aug 25 2005 Simulation Meeting


University of Rochester Geumbong Yu, Yeonsei Chung


## Zmass vs. relative phi

### Monte Carlo

E/p<2.0

Data





Aug 25 2005 Simulation Meeting

University of Rochester Geumbong Yu, Yeonsei Chung